

Personality disorders, schemas, modes in dissociative disorders

Robin van der Linde, Rafaele Huntjens, Nathan Bachrach, Marleen Rijkboer



Introduction

- New conceptualisation of DID → DID mode model
- Shifts between identities are considered shifts between modes

Objectives

To test the DID mode model by investigating:

- 1 personality disorder traits in patients with DID
 - 2 schemas, modes and coping in patients with DID
- in comparison to patients with APD and BPD.


Methods




35 dissociative disorders
138 BPD
37 APD

- ADP-IV
- YSQ-S3
- SMI
- SCI


Results



Schizoid
Schizotypal
Borderline
Avoidant



Disconnection
& rejection



Conclusion

- Severity is at the level of full-blown personality disorders
- Magical thinking
- Hyperassociativity
- Feelings of insufficiency
- Fear of rejection
- Suggestibility
- Para-suicidal behavior
- Potential of applying an adapted form of schema therapy in treating DID

Contactgegevens



Robin van der Linde
r.van.der.linde@rug.nl
GGZ Oost Brabant – Helmond
Rijksuniversiteit Groningen

**CLIËNTEN GEZOCHT VOOR
BEHANDELSTUDIE
“SCHEMATHERAPIE BIJ DE
DISSOCIATIEVE
IDENTITEITSSTOORNIS**

**Kijk vooral ook eens op
www.stbijdis.nl**

